



Total and inorganic As in seafood products caught in an environment facing a mining and industrial area in Sardinia (Italy)

Orletti, Roberta; Sloth, Jens Jørgen; Rasmussen, Rie Romme; Carloni, Cristiano; Griffoni, Francesco; Palombo, Paolo; Velieri, Francesco; Piras, Pierluigi

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Orletti, R., Sloth, J. J., Rasmussen, R. R., Carloni, C., Griffoni, F., Palombo, P., Velieri, F., & Piras, P. (2014). *Total and inorganic As in seafood products caught in an environment facing a mining and industrial area in Sardinia (Italy)*. Poster session presented at 5th International IUPAC Symposium for Trace Elements in Food, Copenhagen, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Total and inorganic As in seafood products caught in an environment facing a mining and industrial area in Sardinia (Italy)

Roberta Orletti¹* (e-mail: r.orletti@izsum.it), Jens J. Sloth², Rie R. Rasmussen², Cristiano Carloni¹,
Francesco Griffoni¹, Paolo Palombo¹, Francesco Velieri¹, Pierluigi Piras³

¹ Istituto Zooprofilattico Sperimentale Umbria – Marche, Via Cupa di Posatora 3, Ancona, Italy
² National Food Institute, Technical University of Denmark, Mørkhøj Bygade 19, DK-2860 Søborg, Denmark
³ Servizio Veterinario IAOA USL 7 Carbonia, Via Dalmazia 831, Carbonia, Italy

INTRODUCTION

The Boi Cerbus lagoon, facing a mining and industrial site in Sardinia (Italy), is an important fishing area for the local population (Fig. 1). Previous studies showed high concentrations of total arsenic (As_{tot}) in fish, molluscs and crustaceans sampled in the lagoon, and a possible exceeding of the JEFCA PTWI value (*Provisional Tolerable Weekly Intake*) by some local consumer groups (teens and children) (1). However, the concentration of inorganic As (As_{inorg}) in the samples should be known for a correct assessment of potential risk, as its toxicity is much higher than that of the organic forms of arsenic.

SAMPLES AND ANALYTICAL METHODS

Eighty samples of 14 different species of fish, molluscs and crustaceans, sampled in the Boi Cerbus lagoon on 3 different seasons (winter, spring and summer 2013), were analysed for As_{tot} by ICP-MS and As_{inorg} by HPLC-ICP-MS (2). The operating scheme is shown in Fig. 2. The HPLC parameters and analytical conditions for ICP-MS are shown in Table 1 and 2, respectively.




Fig. 2. Operating scheme for chemical analysis

Table 1. HPLC analytical conditions.

| Instrument | Perkin Elmer Flexar LC pumps |
|------------------------------|-----------------------------------------------------------------------------|
| Pump | Binary |
| Column | Anion exchange: Transgenomic ICSep ION 120, 120 x 4.6 mm |
| Column flow (ml/min) | 1 |
| Column oven temperature (°C) | 30 |
| Mobile phase | (NH ₄) ₂ CO ₃ 55 mM in 3% MeOH, pH ~ 10.3 |
| Injection volume (µl) | 25 |
| Elution mode | Isocratic |

Table 2. ICP-MS analytical conditions.

| Instrument | Perkin-Elmer SCIEX ELAN 6100DRC II |
|----------------------------|------------------------------------|
| Sample introduction system | |
| Nebulizer | Meinhard |
| Spray chamber | Quartz cyclonic |
| Plasma | |
| RF power (W) | 1300 |
| Carrier Ar flow (L/min) | 0.9-1.1 |
| Mass Spectrometry | |
| Dwell time (ms) | 200 |
| Sweeps/reading | 20 |
| Readings/replicate | 1 |
| Number of replicate | 3 |
| Scanning mode | Peak hopping |
| Monitoring mass | 75 |
| Internal standard | ¹⁰³ Rh (0.1 mg/L) |

RESULTS AND DISCUSSION

Table 3 shows the mean values of As_{tot} and As_{inorg} for each species (+/- SD) and the percentage of As_{inorg}/As_{tot} in bivalves, gastropods, crustaceans and fish. Inorganic forms of As constitute in all cases only a small fraction of total As (max 1% in bivalves). All the data were statistically processed to evaluate significant differences based on season, habitat and *taxon*, in preparation for a subsequent risk assessment. The Kruskal-Wallis test showed no statistically significant differences among the three seasons, nor for the concentrations of total arsenic ($p = 0.2549$), nor for the average percentage of the As_{inorg}/As_{tot} ($p = 0.0817$). The 14 species were divided into two clusters: benthic organisms (Olive green cockle, Banded murex, Mediter. shore crab, Gobies, Combtooth blenny and Common sole) and nektonic organisms (Big-scale sand smelt, Salema, Grey mullets, Gilthead sea bream and European seabass), to evaluate the possible influence of habitats. Using the non-parametric test of Mann-Whitney, the difference between the average levels of As_{tot} in benthic organisms (38.53 ± 5.21 mg/kg) and in nektonic ones (4.89 ± 0.80 mg/kg) was statistically significant ($p < 0.0001$). With regard to average percentage of As_{inorg}/As_{tot} , a not statistically significant difference ($p = 0.1038$) was observed between benthic organisms ($0.21\% \pm 0.04\%$) and nektonic ones ($0.02\% \pm 0.01\%$), using the same nonparametric test. The same species were grouped according to the *taxon* into two new clusters: invertebrates (Olive green cockle, Banded murex, Mediter. shore crab) versus vertebrates (all the other species). According to the Mann-Whitney test, the difference between the average levels of As_{tot} in invertebrates (52.56 ± 7.46 mg/kg) and vertebrates (9.75 ± 1.33 mg/kg) was statistically significant ($p < 0.0001$). Differences due to habitat and *taxon* variability are probably both significant, although currently we are not able to accurately assess the contribution of each one. The difference in mean percentage of As_{inorg}/As_{tot} between invertebrates ($0.31\% \pm 0.06\%$) and vertebrates ($0.03\% \pm 0.01\%$) was statistically significant ($p = 0.0002$). Despite the very high concentrations of As_{tot} detected in some aquatic species, the values of As_{inorg} were in all cases low, even if with different concentrations, due to species-specific characteristics in the metabolism of the element. Therefore a risk assessment based only on the values of As_{tot} can lead to incorrect conclusions.

Table 3. Total and inorganic arsenic mean concentrations (mg/kg w.w. ± S.D.) and their ratio.

| Scientific name | FAO nomenclature | Sample type | N | As_{TOT} | As_{INORG} | As_{INORG}/As_{TOT} % |
|------------------------------------|----------------------|-------------------|----|-----------------|-------------------|-------------------------|
| <i>Cerastoderma glaucum</i> | Olive green cockle | Bivalve mollusc | 11 | 15.5 ± 2.6 | 0.11 ± 0.03 | 0.74 ± 0.22 % |
| <i>Phyllonotus trunculus</i> | Banded murex | Gastropod mollusc | 10 | 106 ± 24 | 0.011 ± 0.012 | 0.01 ± 0.01 % |
| <i>Carcinus aestuarii</i> | Mediter. shore crab | Crustacean | 10 | 40.0 ± 12.6 | 0.054 ± 0.054 | 0.14 ± 0.14 % |
| <i>Gobius niger jozo</i> | Black goby | Fish | 8 | 16.6 ± 4.9 | 0.001 ± 0.001 | <0.01 % |
| <i>Zosterisessor ophiocephalus</i> | Grass goby | Fish | 3 | 34.3 ± 5.7 | 0.004 ± 0.003 | 0.01 ± 0.01 % |
| <i>Solea vulgaris</i> | Common sole | Fish | 6 | 15.4 ± 5.7 | 0.001 | <0.01 % |
| <i>Sparus aurata</i> | Gilthead seabream | Fish | 3 | 15.2 ± 1.9 | 0.001 | 0.00 % |
| <i>Anguilla anguilla</i> | European eel | Fish | 3 | 6.6 ± 5.7 | 0.001 | 0.01 ± 0.01 % |
| <i>Mugil cephalus</i> | Flathead grey mullet | Fish | 6 | 5.2 ± 1.2 | 0.002 ± 0.002 | 0.05 ± 0.04 % |
| <i>Liza aurata</i> | Golden grey mullet | Fish | 6 | 3.7 ± 1.1 | 0.001 ± 0.001 | 0.02 ± 0.02 % |
| <i>Atherina (Hapsetia) boyeri</i> | Big-scale sand smelt | Fish | 4 | 3.0 ± 0.7 | 0.001 | 0.04 ± 0.04 % |
| <i>Salaria basiliscus</i> | Combtooth blenny | Fish | 3 | 2.6 ± 1.7 | 0.007 ± 0.002 | 0.33 ± 0.15 % |
| <i>Dicentrarchus labrax</i> | European seabass | Fish | 5 | 2.0 ± 1.0 | < LOD | <0.01 % |
| <i>Sarpa salpa</i> | Salema | Fish | 2 | 0.49 ± 0.07 | < LOD | <0.01 % |

BIBLIOGRAPHY

- Piras PL, Mulas GF. 2011. Exposure Assessment to food contaminants on edible fish, clams and crustaceans in a coastal environment facing a mining and industrial area in Sardinia (Sulcis-Iglesiente) *Ital J Food Safety* 1: 31-36
- Sloth JJ, Larsen EH, Julshamn K. 2005. Survey of inorganic arsenic in marine animals and marine certified reference materials by anion exchange High-Performance Liquid Chromatography-Inductively Coupled Plasma Mass Spectrometry. *J Agric Food Chem* 53: 6011-6018

This work was funded by the Italian Ministry of Health (project n. IZSUM RC005 2011)




Fig. 1. Boi Cerbus lagoon - Sardinia - Italy